

### EPINEPHRINE DRIP CHART

For a concentration of 4 mcg of epinephrine per milliliter solution.  
1 mg of 1:1,000 mixed in 250 ml of NS

<b>Mix 1mg of epinephrine 1:1,000 in 250ml = 4 mcg/ml</b>	
1 mcg drip = 15 gtt/min	6 mcg drip = 90 gtt/min
2 mcg drip = 30 gtt/min	7 mcg drip = 105 gtt/min
3 mcg drip = 45 gtt/min	8 mcg drip = 120 gtt/min
4 mcg drip = 60 gtt/min	9 mcg drip = 135 gtt/min
5 mcg drip = 75 gtt/min	10 mcg drip = 150 gtt/min

\*Based on a micro drip calibration of 60 drops equal to 1.0 milliliter.

### LIDOCAINE DRIP CHART

For a concentration of 4 mg of lidocaine per milliliter solution.  
1 g mixed in 250 ml of NS.

<b>Mix 1g of lidocaine 1:1,000 in 250ml = 4 mg/ml</b>	
1 mg drip = 15 gtt/min	2 mg drip = 30 gtt/min
3 mg drip = 45 gtt/min	4 mg drip = 60 gtt/min

\*Based on a micro drip calibration of 60 drops equal to 1.0 milliliter.

### INTRANASAL MIDAZOLAM CHART

<b>Patient age</b>	<b>Weight (kg)</b>	<b>Volume in ml (5 mg/ml)</b>	<b>Maximum Dose (mg)</b>
Neonate	3 kg	0.3 ml	0.6 mg
<1 year	6 kg	0.4 ml	1.2 mg
1 year	10 kg	0.5 ml	2.0 mg
2 years	14 kg	0.7 ml	2.8 mg
3 years	16 kg	0.8 ml	3.2 mg
4 years	18 kg	0.9 ml	3.6 mg
5 years	20 kg	1.0 ml	4.0 mg
6 years	22 kg	1.0 ml	4.4 mg
7 years	24 kg	1.1 ml	4.8 mg
8 years	26 kg	1.2 ml	5.2 mg
9 years	28 kg	1.3 ml	5.6 mg
10 years	30 kg	1.4 ml	6.0 mg
11 years	32 kg	1.4 ml	6.4 mg
12 years	34 kg	1.5 ml	6.8 mg
Small teenager	40 kg	1.8 ml	8.0 mg
Adult/large teenager	> 50 kg	2.0 ml	10.0 mg

## PEDIATRIC MEDICATION CHARTS

**DO NOT EXCEED ADULT TOTALS**

**c = concentration**

	Premie	NB	3 Mos.	6 Mos.	1 Yr	2 Yr	4 Yr	6 Yr	8 Yr	10 Yr	12 Yr
<b>Body Length in cm</b>	≤ 53	54-58	59-65	66-74	75-80	81-86	87-99	100-113	114-132	133-158	159-189
<b>Av. Body Wt kg</b>	< 2.5	2.5 - 4	6	7	10	12	16	20	25	34	41
<b>Act Charcoal</b> c=6.25 g/oz. dose = 1 g/kg	1–2.5 g	2.5-4 g	6 g	7 g	10 g	12 g	16 g	20 g	25 g	34 g	41 g
<b>Adenosine</b> c = 3 mg/ml dose = 0.1 mg/kg	-	0.25– 0.4 mg	0.6 mg	0.7 mg	1 mg	1.2 mg	1.6 mg	2.0 mg	2.5 mg	3.4 mg	4.1 mg
<b>Albuterol 1 unit</b> dose (3 ml of 0.083% solution)	2.5 mg	2.5 mg	2.5 mg	2.5 mg	2.5 mg	2.5 mg	2.5 mg	2.5 mg	2.5 mg	2.5 mg	2.5 mg
<b>Amiodarone</b> c= 50 mg/ml dose= 5 mg/kg	-	12.5 - 20 mg	30 mg	35 mg	50 mg	60 mg	80 mg	100 mg	125 mg	170 mg	205 mg
<b>Atropine IV</b> c = 0.1 mg/ml dose = 0.02 mg/kg	-	.1 mg	0.12 mg	0.14 mg	0.2 mg	0.24 mg	0.32 mg	0.4 mg	0.5 mg	0.68 mg	0.82 mg
<b>Dextrose (D10)</b> dose = 5 ml/kg	5–12.5 ml	12.5 - 20 ml	30 ml	35 ml	50 ml	60 ml	80 ml	100 ml	125 ml	170 ml	205 ml
<b>Dextrose (D25)</b> dose = 2 ml/kg	2–5 ml	5 - 8 ml	12 ml	14 ml	20 ml	24 ml	-	-	-	-	-
<b>Dextrose (D50)</b> dose = 1 ml/kg	-	-	-	-	-	-	16 ml	20 ml	25 ml	34 ml	41 ml
<b>Diphenhydramine</b> c = 10 mg/ml dose = 1 mg/kg	1–2.5 mg	2.5 – 4 mg	6 mg	7 mg	10 mg	12 mg	16 mg	20 mg	25 mg	34 mg	41 mg
<b>Epi1:10,000 IV/IO</b> dose = 0.01 mg/kg	0.01- 0.025 mg	0.025 – .04 mg	0.06 mg	0.07 mg	0.1 mg	0.12 mg	0.16 mg	0.2 mg	0.25 mg	0.34 mg	0.41 mg
<b>Epi1:1,000 IM</b> dose = 0.01 mg/kg	-	-	0.06 mg	0.07 mg	0.1 mg	0.12 mg	0.16 mg	0.2 mg	0.25 mg	0.34 mg	0.41 mg

	Premie	NB	3 Mos.	6 Mos.	1 Yr	2 Yr	4 Yr	6 Yr	8 Yr	10 Yr	12 Yr
<b>Body Length in cm</b>	≤ 53	54-58	59-65	66-74	75-80	81-86	87-99	100-113	114-132	133-158	159-189
<b>Av. Body Wt kg</b>	< 2.5	2.5 - 4	6	7	10	12	16	20	25	34	41
<b>Fluid Challenge dose = 20 ml/kg</b>	20–50 ml	50–80 ml	120 ml	140 ml	200 ml	240 ml	320 ml	400 ml	500 ml	680 ml	820 ml
<b>Glucagon c = 1 mg/ml or 1 unit/ml dose = 0.05 mg/kg (up to 1 mg)</b>	0.05 – 0.125 mg	0.125 – 0.2 mg	0.3 mg	0.35 mg	0.5 mg	0.6 mg	0.8 mg	1 mg	1 mg	1 mg	1 mg
<b>Ipratropium bromide c= 0.2 mg/ml dose=</b>	250 mcg	250 mcg	250 mcg	250 mcg	250 mcg	250 mcg	250 mcg	250 mcg	500 mcg	500 mcg	500 mcg
<b>Lidocaine IV c = 20 mg/ml dose = 1 mg/kg</b>	-	2.5 – 4 mg	6 mg	7 mg	10 mg	12 mg	16 mg	20 mg	25 mg	34 mg	41 mg
<b>Magnesium Sulfate c= 500 mg/ml dose= 50 mg/kg</b>	-	0.15-0.2 gm	0.3 gm	0.35 gm	0.5 gm	0.6 gm	0.8 gm	1 gm	1.25 gm	1.7 gm	2.0 gm
<b>Midazolam c = 1 mg/ml dose 0.1 mg/kg</b>	0.2 mg	0.4 mg	0.6 mg	0.7 mg	1 mg	1.2 mg	1.6 mg	2 mg	2.5 mg	3.4 mg	4.1 mg
<b>Morphine c = 10 mg/ml dose = 0.1 mg/kg</b>	-	0.25 - 0.4 mg	0.6 mg	0.7 mg	1 mg	1.2 mg	1.6 mg	2 mg	2.5 mg	3.4 mg	4.1 mg
<b>Naloxone c = 1 mg/ml dose = 0.1 mg/kg</b>	0.1 – 0.25 mg	0.25 - 0.4 mg	0.6 mg	0.7 mg	1 mg	1.2 mg	1.6 mg	2 mg	2 mg	2 mg	2 mg
<b>Sodium Bicarb c = 1 mEq/ml dose = 1 mEq/kg</b>	1 – 2.5 mEq	2.5 – 4 mEq	6 mEq	7 mEq	10 mEq	12 mEq	16 mEq	20 mEq	25 mEq	34 mEq	41 mEq

## TRAUMA TRIAGE CRITERIA

### 1. Physiologic

- Glasgow Coma Score <14;
- Systolic blood pressure (Adult) < 90;
- Systolic blood pressure (Child 7-14) < 85;
- Systolic blood pressure (Child <6) < 70;
- Resp Rate < 10 or > 30 per min;
- Resp Rate Infant <1yo: > 30 per min.

### 2. Anatomic

- Penetrating injuries to head, neck, torso, & extremities proximal to elbow & knee
- Flail chest
- Two or more proximal long bone fractures
- Crushed, degloved, or mangled extremity
- Amputation proximal to wrist & ankle
- Suspected pelvic fracture
- Open or depressed skull fracture
- Traumatic Paralysis

### 3. Mechanism of injury

- Falls (Adult) ≥ 20 ft (1 story = 10 ft)
- Falls ( Child) ≥ 10 ft or 3X child's height
- High Risk Automobile Crash
  - i) Intrusion > 12" at occupant site
  - ii) Ejection from automobile
  - iii) Unrestrained rollover
  - iv) Vehicle telemetry
- Automobile vs. Pedestrian/Bicyclist
  - i) Ped/bicyclist thrown or run over
  - ii) Significant (> 20 mph) impact
- Motorcycle Crash > 20 mph

### 4. Special Considerations

- Older adults: age 55
- Anticoagulation or bleeding disorders
- Burns- Refer to burn triage criteria
- Death in same passenger compartment
- Renal disease requiring dialysis
- Pregnancy > 20 weeks with complaint of injury
- EMS provider judgment

### Vision, Aphasia, Neglect (VAN) Scale

	Normal	Mild	Moderate	Severe
How weak is the Patient? Raise both arms.	No signs of weakness	Minor Drift	Severe drift- touches or nearly touches the ground	Flaccid or no antigravity

Visual Disturbances	Aphasia	Neglect
Field cut (which side) (4 quadrants)	Expressive (inability to speak or paraphasic errors); do not count slurred words (repeat & name 2 objects)	Forced gaze or inability to track to one side.
Double Vision (ask pt. to look to right then left; evaluate for uneven eyes)	Receptive (not understanding or following commands) i.e., close eyes, make a fist.	Unable to feel both sides at the same time, or unable to identify own arm.
Blind- New Onset	Mixed	Ignoring one side.
None	None	None

\*Patient must have weakness and one or more of the V, A, N to be VAN Positive.

### Sepsis Screen

Patient should be presumed to be septic if the patient meets two or more of the following criteria, with no other identifiable cause;

1. Temperature > 100.4° or < 96°
2. Heart rate > 90
3. Respiratory rate > 20

**TUOLUMNE COUNTY EMS AGENCY  
FIBRINOLYTIC CHECKLIST**

Date \_\_\_\_\_ PCR# \_\_\_\_\_ Receiving Facility \_\_\_\_\_

Patient Name \_\_\_\_\_ DOB \_\_\_\_\_ Medic Unit \_\_\_\_\_

If ANY of the following is checked YES, fibrinolysis MAY be contraindicated:

- |   | YES | NO  |
|---|-----|-----|
| 1. Chest pain lasting greater than 12 hours                                   | ( ) | ( ) |
| 2. Systolic BP greater than 180 mmHg  | ( ) | ( ) |
| 3. Diastolic BP greater than 100 mmHg   | ( ) | ( ) |
| 4. Age younger than 35 if male or 40 if female                                | ( ) | ( ) |
| 5. History of stroke, TIA, brain tumor, A-V malformation or other CNS disease | ( ) | ( ) |
| 6. Internal bleeding in past 2-4 weeks  | ( ) | ( ) |
| 7. Surgery or trauma in past 6 weeks, including laser eye surgery             | ( ) | ( ) |
| 8. Closed head/facial trauma past 3 months                                    | ( ) | ( ) |
| 9. Bleeding or clotting problems or on anticoagulants                         | ( ) | ( ) |
| 10. Pregnant female   | ( ) | ( ) |
| 11. Terminal illness  | ( ) | ( ) |
| 12. Serious systemic disease, including liver or kidney disease               | ( ) | ( ) |
| 13. Previous hypersensitivity to reteplase                                    | ( ) | ( ) |

High Risk?


If any of the following are checked YES, consider transport to PCI facility:

- |   | YES | NO  |
|---|-----|-----|
| 1. Heart rate greater than or equal to 100 bpm AND SBP less than 100 mmHg | ( ) | ( ) |
| 2. Pulmonary edema (rales)  | ( ) | ( ) |
| 3. Signs of shock (cool, clammy)  | ( ) | ( ) |
| 4. Contraindications to fibrinolytic therapy                              | ( ) | ( ) |

\_\_\_\_\_  
*Assessing Paramedic Signature*

\_\_\_\_\_  
*Paramedic License #*

### Burn Surface Area



Rate based on a single micro infusion set, with drip calibration of 60 drops equal to 1.0 milliliter.

Rate based on a single macro infusion set drip, with calibration of 10 drops equal to 1.0 milliliter.

Rate based on two macro infusion sets, with drip calibration of 10 drops equal to 1.0 milliliter. Both IVs should be infusing at the prescribed drip rate.

#### Drip Rate Based on Surface Area Burned (2° & 3°) and Body Weight

Wt kg	5	10	15	20	25	30	35	40	45	50	55	60	65	70	75
9%	15	30	45	45	60	75	90	100	100	100	20	25	25	30	30
9.5%	15	30	45	45	60	75	90	100	100	100	20	25	25	30	30
10%	15	30	45	45	100	100	90	100	100	20	25	25	30	30	30
15%	15	45	60	75	100	100	25	25	30	30	30	45	45	45	45
17%	30	45	60	90	100	20	25	30	30	45	45	45	45	45	60
18%	30	45	60	90	100	25	30	30	30	45	45	45	60	60	60
19%	30	45	75	100	100	25	30	30	30	45	45	45	60	60	60
24%	30	45	90	20	25	30	30	45	45	45	45	60	60	75	75
24.5%	30	45	90	20	25	30	30	45	45	45	60	60	60	75	75
28%	45	75	100	25	30	45	45	45	60	60	60	75	75	75	90
30%	45	75	100	25	30	45	45	45	60	60	60	75	75	90	90
32%	45	75	20	30	30	45	45	45	60	60	75	75	90	90	100
36%	45	90	25	30	45	45	60	60	60	75	75	90	90	100	100
37%	45	90	25	30	45	45	60	60	60	75	75	90	100	100	100
38%	45	100	25	30	45	45	60	60	75	75	75	90	100	100	100
39%	45	100	25	30	45	45	60	60	75	75	90	90	100	100	100
41.5%	45	100	25	45	45	60	60	75	75	90	90	100	100	60	60
45%	60	100	30	45	45	60	60	75	75	90	100	100	60	60	60
46%	60	100	30	45	45	60	60	75	75	90	100	100	60	60	60
47%	60	100	30	45	45	60	75	75	90	90	100	100	60	60	60
49%	60	20	30	45	45	60	75	75	90	100	100	60	60	60	75
51%	60	20	30	45	45	60	75	75	90	100	100	60	60	60	75
54%	60	25	30	45	60	60	75	90	100	100	60	60	60	75	75
54.5%	60	25	30	45	60	75	75	90	100	100	60	60	60	75	75
56.5%	75	25	45	45	60	75	75	90	100	100	60	60	75	75	90
58.5%	75	25	45	45	60	75	75	90	100	60	60	60	75	75	90
62%	75	25	45	45	60	75	90	100	100	60	60	75	75	90	90
63%	75	30	45	45	60	75	90	100	100	60	60	75	75	90	100
64%	75	30	45	45	60	75	90	100	60	60	60	75	75	100	100
66%	90	30	45	45	75	75	90	100	60	60	75	75	90	100	100
68%	90	30	45	60	75	75	100	100	60	60	75	75	90	100	100
71%	90	30	45	60	75	90	100	100	60	60	75	90	90	100	100
71.5%	90	30	45	60	75	90	100	100	60	75	90	90	90	100	100
75.5%	100	30	45	60	75	90	100	60	60	75	90	90	100	100	100
81%	100	30	45	60	75	100	100	60	75	75	90	100	100	100	120
82%	100	30	45	75	90	100	100	60	75	75	90	100	100	100	120
90.5%	100	45	60	75	90	100	60	75	75	90	100	100	100	120	150
91%	100	45	60	75	90	100	60	75	75	90	100	100	100	120	150
100%	20	45	60	90	100	60	60	75	90	100	100	100	120	150	150

**Drip Rate Based on Surface Area Burned (2° & 3°) and Body Weight**

<b>Wt kg</b>	<b>80</b>	<b>85</b>	<b>90</b>	<b>95</b>	<b>100</b>	<b>105</b>	<b>110</b>	<b>115</b>	<b>120</b>	<b>125</b>	<b>130</b>	<b>135</b>	<b>140</b>	<b>145</b>	<b>150</b>
<b>9%</b>	30	45	45	45	45	45	45	45	45	45	45	45	60	60	60
<b>9.5%</b>	30	45	45	45	45	45	45	45	45	45	45	45	60	60	60
<b>10%</b>	30	45	45	45	45	45	45	45	45	45	45	60	60	60	60
<b>15%</b>	45	45	60	60	60	60	75	75	75	75	75	75	90	90	90
<b>17%</b>	60	60	60	60	75	75	75	75	75	90	90	90	100	100	100
<b>18%</b>	60	60	60	75	75	75	75	75	90	90	100	100	100	100	100
<b>19%</b>	60	60	75	75	75	75	90	90	90	90	100	100	100	100	100
<b>24%</b>	75	90	90	90	100	100	100	100	60	60	60	60	60	60	75
<b>24.5%</b>	75	90	90	90	100	100	100	100	60	60	60	60	60	60	75
<b>28%</b>	90	100	100	100	100	100	100	100	60	75	75	75	75	75	75
<b>30%</b>	100	100	100	100	60	60	60	60	75	75	75	75	75	90	90
<b>32%</b>	100	100	60	60	60	60	60	75	75	75	75	90	90	100	100
<b>36%</b>	60	60	60	60	75	75	75	75	90	90	90	100	100	100	100
<b>37%</b>	60	60	60	60	75	75	75	90	90	100	100	100	100	100	100
<b>38%</b>	60	60	60	75	75	75	75	90	90	100	100	100	100	100	100
<b>39%</b>	60	60	60	75	75	75	90	90	100	100	100	100	100	100	100
<b>41.5%</b>	60	60	75	75	75	90	90	100	100	100	100	100	100	120	120
<b>45%</b>	75	75	75	90	90	100	100	100	100	100	100	120	120	120	150
<b>46%</b>	75	75	75	90	100	100	100	100	100	100	120	120	120	120	150
<b>47%</b>	75	75	75	90	100	100	100	100	100	100	120	120	120	150	150
<b>49%</b>	75	75	90	100	100	100	100	100	100	120	120	120	150	150	150
<b>51%</b>	75	90	100	100	100	100	100	100	120	120	120	150	150	150	150
<b>54%</b>	90	100	100	100	100	100	100	120	120	150	150	150	150	150	150
<b>54.5%</b>	90	100	100	100	100	100	120	120	120	150	150	150	150	150	150
<b>56.5%</b>	90	100	100	100	100	100	120	120	150	150	150	150	150	150	150
<b>58.5%</b>	100	100	100	100	100	120	120	120	150	150	150	150	150	150	150
<b>62%</b>	100	100	100	100	120	120	150	150	150	150	150	150	150	150	150
<b>63%</b>	100	100	100	120	120	120	150	150	150	150	150	150	150	150	150
<b>64%</b>	100	100	100	120	120	120	150	150	150	150	150	150	150	150	WO
<b>66%</b>	100	100	100	120	120	150	150	150	150	150	150	150	150	150	WO
<b>68%</b>	100	100	120	120	150	150	150	150	150	150	150	150	150	WO	WO
<b>71%</b>	100	120	120	120	150	150	150	150	150	150	150	WO	WO	WO	WO
<b>71.5%</b>	100	120	120	150	150	150	150	150	150	150	150	WO	WO	WO	WO
<b>75.5%</b>	120	120	150	150	150	150	150	150	150	150	WO	WO	WO	WO	WO
<b>81%</b>	120	150	150	150	150	150	150	150	WO	WO	WO	WO	WO	WO	WO
<b>82%</b>	120	150	150	150	150	150	150	150	WO	WO	WO	WO	WO	WO	WO
<b>90.5%</b>	150	150	150	150	150	150	WO	WO	WO	WO	WO	WO	WO	WO	WO
<b>91%</b>	150	150	150	150	150	150	WO	WO	WO	WO	WO	WO	WO	WO	WO
<b>100%</b>	150	150	150	150	WO	WO	WO	WO	WO	WO	WO	WO	WO	WO	WO

**ABBREVIATIONS**

ALS:	Advanced Life Support
BLS:	Basic Life Support
BVM:	Bag Valve Mask
CHF:	Congestive Heart Failure
CNS:	Central Nervous System
CPAP:	Continuous Positive Airway Pressure
CWI:	Cold Water Immersion
D10:	Dextrose 10%
dL:	deciliter
ETI:	Endotracheal Intubation
GCS:	Glasgow Coma Score
G:	Grams
HP-CPR:	High performance Cardiopulmonary Resuscitation
IM:	Intramuscular
IN:	Intranasal
IO:	Intraosseous
IV:	Intravenous
IVP:	Intravenous Push
J:	Joules
kg:	kilograms
mcg:	micrograms
mEq:	milliequivalent
mg:	milligrams
mL:	milliliters
NG:	Nasogastric
NS:	Normal Saline
NTG:	Nitroglycerin
ROSC:	Return of Spontaneous Circulation
ODT:	Oral Disintegrating Tablet
SBP:	Systolic Blood Pressure
SGA:	Supraglottic Airway
TCP:	Transcutaneous Pacing