

TENSION PNEUMOTHORAX (A13)

Physical signs may include: decreased breath sounds, increased resonance on the side of collapsed lung, tracheal deviation, asymmetrical chest motion and crepitus.

	F	E	O	P	D
ASSESSMENT	X	X	X	X	
BLS AIRWAY: okay if airway patent. Support ventilations with appropriate airway adjuncts. Ventilate with 100% oxygen.	X	X	X	X	
PULSE OXIMETRY: apply and monitor.		X	X	X	
CAPNOGRAPHY: apply and monitor.				X	
OXYGEN: if pulse oximetry < 94% or signs of respiratory distress or hypoperfusion.	X	X	X	X	
ECG MONITOR: lead placement may be delegated. Treat as indicated.				X	
NEEDLE THORACOSTOMY: on affected side(s) between 2 nd & 3 rd intercostal space midclavicular line, OR between 4 th & 5 th intercostal space midaxillary line. Place catheter just above the rib to avoid the intercostal artery. Place approved chest seal over or one-way valve on the catheter. Repeat if suspected catheter occlusion.				X	
ADVANCED AIRWAY: if GCS is < 8 and not rapidly improving, consider SGA or ETI.				X	
VASCULAR ACCESS: IV/IO, rate as indicated.				X	
ASSESSMENT: continue to monitor for signs of recurrence of tension pneumothorax.	X	X	X	X	

EFFECTIVE: November 1, 2023

Provider Key: F = First Responder/EMR
P = Paramedic

E = EMT O = EMT Local Optional SOP
D = Base Hospital Physician Order Required